

---

# Guidance on Air Medical Transport for Patients with Ebola Virus Disease

---

## On this Page

- Background
- A. Air Medical Transport of Patients with EVD: General Considerations
- B. Patient Placement
- C. Infection Control
- D. Patients Requiring Special Consideration
- E. Waste Disposal
- F. Cleaning and Disinfection
- G. Transport of an Asymptomatic Exposed Person
- H. Logistical Planning and Post-Mission Follow-Up
- I. Ground/In-Flight Emergency Procedures
- Related Pages

**Who this is for:** Operators of air medical transport (AMT) services that are considering transport of patients with Ebola virus disease (EVD) and healthcare providers who will be onboard. This guidance applies to AMT flights of any duration and using any type of aircraft. **The guidance does not apply to commercial passenger aircraft or other types of patient transport vehicles (for example, ground ambulances).**

**What this is for:** Guidance for safe transportation of patients with EVD by AMT.

**How to use:** Use this guidance to ensure safety of healthcare personnel and patients during AMT of patients with EVD. Explain to pilots, other aircraft personnel, and cleaning crews what special actions should occur before, during and after transport, and how to keep themselves safe.

## Key Points

- Transport must be coordinated with public health and civil aviation authorities at origin and destination.
- Infection control policies and procedures should be established before and implemented during all phases of patient transport.
- A portable isolation unit is recommended to contain infected materials and minimize contamination of

the aircraft.

- Personnel providing care during transport should be trained in clinical management, infection control, and correct use of personal protective equipment (PPE).
- PPE should be used by all those in the patient care area or who may have contact with patients or their body fluids; infection control guidelines should be followed, and procedures that could increase the risk of exposure to the patient's body fluids should be avoided.

## Background

Air medical transport (AMT) is a unique patient care setting. Unlike ground medical transport, such as in a ground ambulance, the aircraft usually does not stop or re-supply during transport and the mission usually exceeds several hours. During AMT, a patient's condition can deteriorate, needing additional interventions that could result in an increased risk of exposure for healthcare providers. The recommendations in this guidance are based on standard infection prevention and control practices for Ebola and Marburg hemorrhagic fever, AMT standards, aircraft-specific safety considerations, and established international notification protocols.

Ebola virus is transmitted in healthcare settings by direct contact with infectious blood or body fluids (for example, blood, urine, feces, or vomit); accidental injuries by sharps, such as scalpel blades or needles, that are contaminated with infectious material; and splashes to unprotected mucous membranes of the eyes, nose, or mouth. Infection control measures for patients with Ebola virus disease (EVD) include standard, contact, and droplet precautions. Injection therapy, blood sampling, and other procedures that require the use of needles and other sharp implements should be limited to what is essential for patient care. Procedures likely to generate splashing (such as transferring liquid waste from one container to another) should be performed only when necessary and with careful adherence to correct use of personal protective equipment (PPE). Procedures that might generate aerosols, such as intubation, should be avoided unless necessary for patient care. If it is necessary to perform such a procedure, healthcare providers should wear appropriately fit-tested respirators that provide at least 95% filtering efficiency (such as a NIOSH-certified N-95 or higher-level filtering facepiece respirator).

The decision to transport a patient with EVD should be carefully considered and thoroughly planned in discussion with relevant public health agencies, civil aviation authorities, and personnel from the receiving facility. Factors that should weigh into the decision include whether the patient is clinically stable enough to travel (i.e., whether the patient's condition could deteriorate during transport), and whether personnel providing care during transport are trained in clinical management, infection control, and correct use of PPE.

[^ Top of Page](#)

### A. Air Medical Transport of Patients with EVD: General Considerations

- Patients with EVD should be transported on a dedicated AMT mission.
  - **Other patients who do not have laboratory-confirmed EVD should not be onboard.**

- Only people directly involved in patient care or operating the aircraft should be onboard. If people other than those caring for the patient and operating the aircraft, such as a patient's family member, need to be on the aircraft, decisions should be made on a case-by-case basis in consultation with public health authorities and, as necessary, aviation authorities.
- Pilots, other necessary crew members, and ground crew should follow routine procedures for AMT missions. Anyone not involved in direct patient care should remain at least 3 feet (1 meter) away from the patient during movement of the patient onto and off the aircraft, and should not enter the designated patient care area.
- If available, a portable isolation unit (see [C. Infection Control](#) below) should be used to contain infectious materials. Use of a portable isolation unit minimizes the need to clean and decontaminate the aircraft after the mission.
- Infection control involves a variety of precautionary measures (see [C. Infection Control](#) for details):
  - Eliminate exposure to body fluids.
  - Minimize contamination of the aircraft. Disinfect surfaces that are soiled during patient care promptly (see [F. Cleaning and Disinfection](#) below).
  - Use personal protective equipment (PPE) correctly.
  - Avoid procedures that require the use of sharps or that could create splashes of infectious material.
- Plan that the crew will need to take breaks in compliance with crew rest requirements.
- Coordinate international transport of patients with EVD with public health and civil aviation authorities at origin and destination.
- In the United States, EVD is a disease for which [federal isolation and quarantine](#) are authorized.
  - All transport of patients with EVD destined for a U.S. airport must be reported to CDC before arrival, preferably before travel, by calling the CDC Emergency Operations Center (EOC) at **+1 770-488-7100** (24-hour number). U.S. federal regulations require the commander of an aircraft destined for the United States to report to CDC any deaths and certain illnesses among travelers before arrival. For more information, see [Guidance for Airlines on Reporting Onboard Deaths or Illnesses to CDC](#).
  - Reports from flights operating in U.S. airspace provided to the Federal Aviation Administration Air Traffic Services (ATS) units are shared with the CDC EOC.
- International movement of patients with EVD might additionally require special approvals by aircraft-servicing or fueling, or patient rest-stop locations, and countries that will be over-flown, if applicable.
- If the AMT experiences an in-flight incident (such as a change to the patient's condition requiring the flight to divert or be supported by special handling), the pilot should advise the appropriate ATS unit, to facilitate coordination with the responsible public health authorities and the implementation of any needed air traffic management related support. The pilot also should follow applicable company procedures.
  - For more information, see the International Civil Aviation Organization's (ICAO) Annex 9, Facilitation, and Document 4444, Procedures for Air Navigation Services – Air Traffic Management.

## B. Patient Placement

- If transport of more than one patient with confirmed EVD is planned, the patients can be isolated together as a group.

^ [Top of Page](#)

## C. Infection Control

### 1. Designation of an “isolation area”

- Use of a disposable, portable isolation unit is recommended to contain infectious waste (such as soiled absorbent pads, emesis [vomit] basins, portable toilet) and to prevent contamination of the aircraft cabin. After use, the unit should be discarded (see [E. Waste Disposal](#) below). If an isolation unit is not available, a perimeter should be established for designating “clean” and “dirty” areas that identifies where gowns and gloves should be put on and removed. The distance will depend on the area required for patient care support as well as designated space for safe removal of PPE. A minimum distance of 6 feet (2 meters) from the patient is recommended. Surfaces should be smooth, nonporous, and fluid-impermeable to allow appropriate cleaning and disinfection.
- Materials required for patient care, including PPE, should be stored outside of the isolation area.
- Containers for soiled linen, waste, and reusable equipment should be placed inside the isolation area. Supplies for cleaning spills should be kept in the area, including appropriate disinfectants (see [F. Cleaning and Disinfection](#) below).
- A portable chemical toilet dedicated for patient use should be placed within the isolation area. The lid of the toilet should be kept closed.

Personnel who are within the isolation unit or designated isolation area should wear PPE as described in the CDC guidance: [Guidance on Personal Protective Equipment To Be Used by Healthcare Workers During Management of Patients with Ebola Virus Disease in U.S. Hospitals, Including Procedures for Putting On \(Donning\) and Removing \(Doffing\)](#). Outside of the isolation unit or designated isolation area, PPE is not needed

### 2. Minimize opportunities for exposure

- Limit use of needles and other sharps as much as possible.
- If use of needles or other sharps is necessary for patient care, handle these items with extreme care and discard them immediately after use in closed, puncture-proof containers.
- Use disposable equipment whenever possible.
- Avoid aerosol-generating procedures.
- Avoid procedures that can generate splashes of infectious material, such as transferring liquid waste from one container to another.

### 3. Personal protective equipment and procedures

- For recommended personal protective equipment (PPE) and procedures for putting on and removing PPE, refer to [Guidance on Personal Protective Equipment To Be Used by Healthcare Workers During Management of Patients with Ebola Virus Disease in U.S. Hospitals, Including Procedures for Putting On \(Donning\) and Removing \(Doffing\)](#).

^ [Top of Page](#)

## D. Patients Requiring Special Consideration

- Oxygen support may be provided during flight with simple and non-rebreather face masks.
- Mechanical ventilators for patients with EVD should provide HEPA or equivalent filtration of airflow exhaust.
- AMT services should consult their ventilator equipment manufacturer to confirm appropriate filtration capability and the effect of filtration on positive-pressure ventilation.

^ [Top of Page](#)

## E. Waste Disposal

### 1. Management of waste during AMT

- Collect dry solid waste (such as used gloves, dressings) in leakproof biohazard bags.
- Collect waste soaked with blood or body fluids in leakproof biohazard bags or containers.
- Discard sharp items, such as used needles or scalpel blades, immediately after use in puncture-proof sharps containers.
- Store suctioned fluids and secretions in sealed containers. Handling patient body fluids might create splashes and should be avoided.
- If bedpans or urinals are needed, excretions, such as feces or urine, may be carefully poured down the dedicated portable patient toilet, taking care to avoid splashes and spills. If splashing or spills cannot be avoided, fluids should be kept in a sealed container until they can be disposed of safely. If necessary, the entire bedpan may be placed in a sealed container.

### 2. Disposal of waste after mission completion

- In the United States, Ebola virus is a Category A infectious substance regulated by the U.S. Department of Transportation's (DOT) Hazardous Materials Regulations (HMR).
  - Any item transported for disposal that is contaminated with a Category A infectious substance must be packaged and transported in accordance with DOT regulations found in [49 Code of Federal Regulations Parts 171-180 \(http://www.ecfr.gov/cgi-bin/text-idx?SID=5ca0292b2d30bf3287b17309e7b540e0&tpl=/ecfrbrowse/Title49/49CISubchapC.tpl\)](#). This includes used medical equipment or PPE, contaminated linens, and other medical waste collected

during transport of a patient with confirmed or suspected EVD.

- All applicable requirements of the HMR can be found on the DOT website:

<http://phmsa.dot.gov/hazmat> (<http://phmsa.dot.gov/hazmat>) or in DOT Guidance for Transporting Ebola-Contaminated items

(<http://phmsa.dot.gov/portal/site/PHMSA/menuitem.6f23687cf7b00b0f22e4c6962d9c8789/?vgnextoid=4d1800e36b978410VgnVCM100000d2c97898RCRD&vgnnextchannel=d248724dd7d6c010VgnVCM10000080e8a8c0RCRD&vgnnext>).

- For questions on hazardous materials packaging and transportation regulations, contact the U.S. DOT HazMat Information Center at 800-467-4922.
- Outside of the United States, follow applicable regulations for waste disposal.
- Plans for waste disposal should be discussed in advance with waste handlers at the airport of arrival and the destination medical facility.

^ [Top of Page](#)

## F. Cleaning and Disinfection

- Recommended PPE should be worn by personnel responsible for cleaning environmental surfaces and should, at a minimum, be consistent with standard, contact, and droplet precautions.
- Environmental surfaces should be disinfected using a U.S. Environmental Protection Agency (EPA)-registered hospital disinfectant for use against the Ebola virus that is approved by the aircraft manufacturer. Currently, no EPA-registered hospital disinfectant products will have a statement on the label that specifically says it can kill Ebola virus. Use an Environmental Protection Agency (EPA)-registered hospital disinfectants with label claims for hospital disinfection (or the equivalent microbial pathogen claims) and claims against non-enveloped viruses (e.g., norovirus, rotavirus, adenovirus, poliovirus). These disinfectants should be used in accordance with the manufacturer's labeled instructions. One simple way to identify an appropriate product effective against Ebola virus is to use a product included in EPA's List L: Disinfectants for Use Against the Ebola Virus (<http://www.epa.gov/oppad001/list-l-ebola-virus.html>). Disinfectant should be available in spray bottles or as commercially prepared wipes for use during transport.
- Any surface that becomes potentially contaminated during transport immediately should be sprayed and wiped clean, or simply wiped clean using a commercially prepared disinfectant wipe.
- A blood spill or spill of other body fluid or substance (for example, urine, feces, or vomit) should be managed through removal of bulk spill matter, cleaning the site, and then disinfecting the site.
  - For large spills, a chemical disinfectant (EPA-registered hospital disinfectant) with sufficient potency is needed to overcome the tendency of proteins in blood and other body substances to neutralize the disinfectant's active ingredient.
  - An EPA-registered hospital disinfectant with label claims for hospital disinfection (or the equivalent microbial pathogen claims) and claims against non-enveloped viruses (e.g., norovirus, rotavirus,

adenovirus, poliovirus) should be used. One simple way to identify an appropriate product effective against Ebola virus is to use a product included in EPA's List L: [Disinfectants for Use Against the Ebola Virus \(http://www.epa.gov/oppad001/list-l-ebola-virus.html\)](http://www.epa.gov/oppad001/list-l-ebola-virus.html).

- Instructions for cleaning and decontaminating surfaces or objects soiled with blood or body fluids should be followed to treat the spill before covering with absorbent material and wiping up.
- After the bulk waste is wiped up, the surface should be disinfected as described above.
- All patient-care areas (including stretchers, railings, medical equipment control panels, and adjacent flooring, walls, and work surfaces likely to be directly contaminated during care) should be considered contaminated and therefore cleaned and disinfected after AMT.
- Compressed air or pressurized water should not be used for cleaning the aircraft.
- Areas of the aircraft that were not used for patient care should be cleaned and maintained as per routine practices according to manufacturers' recommendations.
- Contaminated reusable patient care equipment should be placed in biohazard bags and labeled for cleaning and disinfection at the AMT service medical equipment section.
- Reusable equipment should be cleaned and disinfected, according to manufacturer's instructions, by trained personnel wearing correct PPE.
- Use of reusable items with porous surfaces that cannot be easily replaced should be avoided.
- Only mattresses and pillows with plastic or other fluid-impermeable covering should be used. Coverings should be disinfected after use or discarded (see [E. Waste Disposal](#)).
- To prevent exposure among laundry staff, all contaminated linens should be discarded.
- Any contaminated nonfluid-impermeable items, such as web seats, seat cushions, pillows, or mattresses should be discarded.
- After completing cleaning tasks, including cleaning and disinfection of reusable equipment, cleaning personnel should carefully remove and dispose of PPE as described in CDC's guidance: [Guidance on Personal Protective Equipment To Be Used by Healthcare Workers During Management of Patients with Ebola Virus Disease in U.S. Hospitals, Including Procedures for Putting On \(Donning\) and Removing \(Doffing\)](#). They should [perform \(http://www.cdc.gov/handhygiene/\)](http://www.cdc.gov/handhygiene/) hand hygiene with soap and water or an alcohol-based hand sanitizer. Soap and water should be used if hands are visibly soiled.

^ [Top of Page](#)

## G. Transport of an Asymptomatic Exposed Person

Persons who have had [exposure to Ebola virus](#) should not travel by commercial aircraft because of the risk of developing symptoms in transit and potentially exposing others onboard. When transporting someone who has been exposed but who does not have symptoms, infection control precautions and use of PPE are not required. However, PPE should be available in case it is needed. The person who has been exposed should be monitored carefully, including temperature measurements every 12 hours, for elevated body temperature or subjective fever or [other signs or symptoms compatible with EVD](#), including severe headache, fatigue, muscle

pain, vomiting, diarrhea, abdominal pain, or unexplained hemorrhage. The exposed person immediately should notify transport personnel if he or she suspects fever or develops other symptoms. If symptoms develop, standard, contact, and droplet precautions should be implemented immediately.

^ [Top of Page](#)

## H. Logistical Planning and Post-Mission Follow-Up

- Sufficient infection control supplies should be onboard to support the expected duration of the mission plus additional time in the event the aircraft experiences maintenance delays or weather diversions.
- Flight planning should identify diversion airfields and coordinate with authorities responsible for those locations in advance. The potential need for coordination with ATS units and public health authorities responsible for those countries being overflown also should be taken into consideration.
- Once the mission is completed, the AMT team should provide the following information to their medical director: mission number/date; address of the team/aircraft basing; duration of patient transport; contact information for all team members; crew positions (including estimated duration of direct patient care provided); and a description of any recognized breaches in infection control.
- Personnel with any potential exposures to infectious material during the mission should be evaluated immediately in consultation with public health authorities so a plan can be implemented for appropriate monitoring and medical evaluation and testing if indicated. Additional [monitoring and movement restrictions](#) may be imposed by public health authorities for personnel with unprotected exposure to a patient with EVD.
- All mission personnel should be advised to self-monitor for a period of 21 days after the last known contact with the patient with EVD. They should immediately report elevated body temperature or subjective fever or any other [signs or symptoms consistent with EVD](#) to their supervisor.
- AMT services should designate people responsible for performing post-mission monitoring of mission personnel and reporting results to the AMT service medical director.

^ [Top of Page](#)

## I. Ground/In-Flight Emergency Procedures

AMT service providers should have a written plan addressing patient handling during in-flight or ground emergency situations. Certain activities, such as putting on life vests and rapidly evacuating the patient from the aircraft, may create special exposure risks. Use of PPE must be weighed against time constraints and the nature of the emergency conditions (such as smoke in the cabin, sudden cabin decompression).

^ [Top of Page](#)

---

Page last reviewed: December 12, 2014

Page last updated: December 12, 2014

Content source: Centers for Disease Control and Prevention (/index.htm)

National Center for Emerging and Zoonotic Infectious Diseases (NCEZID) (/ncezid/index.html)



[Division of High-Consequence Pathogens and Pathology \(DHCPP\) \(/ncezid/dhcpp/index.html\)](/ncezid/dhcpp/index.html)

[Viral Special Pathogens Branch \(VSPB\) \(/ncezid/dhcpp/vspb/index.html\)](/ncezid/dhcpp/vspb/index.html)